WO 2005/054444 PCT/KR2004/003024

What is claimed is:

5

10

15

1. A chemically defined medium for fermentation culture of a strain of the genus *Candida*, which comprises 1-10 g/ ℓ of urea, 1-10 g/ ℓ of potassium diphosphate, 0.01-1 g/ ℓ of magnesium sulfate, 0.1-10 mg/ ℓ of MnSO₄ · 4H₂O, 0.1-10 mg/ ℓ of CoCl₂ · 6H₂O, 0.1-10 mg/ ℓ of NaMoO₄ · 2H₂O, 0.1-10 mg/ ℓ of ZnSO₄ · 7H₂O, 0.1-10 mg/ ℓ of AlCl₃ · 6H₂O, 0.1-10 mg/ ℓ of CuCl₂ · 2H₂O, 0.01-5 mg/ ℓ of H₃BO₃, 1-100 mg/ ℓ of FeSO₄ · 7H₂O, 0.1-10 mg/ ℓ of ascorbic acid, 1-100 mg/ ℓ of biotin, 1-100 mg/ ℓ of choline, and 0.1-10 mg/ ℓ of pyridoxine.

2. A process for producing xylitol in high yield by recycling culture of a strain of the genus *Candida*, which comprises the steps of:

inoculating the strain in a xylose-containing medium and culturing the strain in the xylose-containing medium in a bioreactor;

releasing a culture from the bioreactor and introducing a fresh xylosecontaining medium to the bioreactor continuously;

separating the strain and a culture filtrate from the culture; and

recycling the strain to the bioreactor and recovering xylitol from the culture filtrate.

- 3. The process of claim 2, wherein the strain of the genus *Candida* is *Candida tropicalis* or its mutant strain.
- 4. The process of claim 2, wherein the xylose-containing medium is the chemically defined medium of claim 1 or a complex medium.

20

WO 2005/054444 PCT/KR2004/003024

5. The process of claim 2, wherein the culturing is performed by a fedbatch culture or a batch culture.

- 6. The process of claim 5, wherein in the fed-batch culture, the medium is gradually supplemented with xylose so that the concentration of xylose is maintained at $40-50g/\ell$ on the basis of the medium.
 - 7. The process of any one of claims 2, 4, 5, and 6, wherein the culturing is performed at an agitation speed of 400-600rpm.
 - 8. The process of claim 2, wherein the separation of the strain and the culture filtrate from the culture is performed by a vacuum microfiltration system or a centrifuge.

10

15 9. The process of claim 2 or 8, wherein the separated strain is concentrated to a density of $10-100g/\ell$ and recycled.